

# Update

## Neural Network based Jet Charge Tagger

G. Barker, M. Feindt, U. Kerzel, C. Lecci  
G. Bauer, A. Belloni, I. Kravchenko,  
N. Leonardo, S. Menzemer, J. Miles, Ch. Paus, A. Rakitin,  
G. Gomez-Ceballos, J. Piedra, A. Ruiz, I. Vila,  
I. Furić

Karlsruhe, MIT, Cantabria, U. Chicago

CP Violation and Mixing Meeting, March 1<sup>st</sup>, 2005

## Update

Since the pre-blessing talk (02/17)

- bug fix for  $Q_{jet}$  computation  $\rightarrow \epsilon D^2$  slightly changed
- computed NNJQT performance with exclusion of SLT tagged events
- considered track probability optimization on three samples
  - $\rightarrow$  tracks with L00 hits
  - $\rightarrow$  SI tracks without L00 hits
  - $\rightarrow$  **COT only tracks**

## Updated $\epsilon D^2$ numbers

$Q_{jet}$  computed with no separating tracks with and without L0 hits **FIXED**

Old numbers were...

	$\epsilon, \%$	effective $D, \%$	$\epsilon D^2$
$e$ +SVT combined	$95.47 \pm 0.15$	$9.77 \pm 0.18$	<b><math>0.912 \pm 0.034</math></b>
$\mu$ +SVT combined	$95.73 \pm 0.12$	$9.45 \pm 0.16$	<b><math>0.855 \pm 0.029</math></b>

New numbers are...

$e$ +SVT combined	$95.48 \pm 0.15$	$9.80 \pm 0.16$	<b><math>0.917 \pm 0.031</math></b>
$\mu$ +SVT combined	$95.72 \pm 0.12$	$9.90 \pm 0.15$	<b><math>0.938 \pm 0.029</math></b>

## Soft Lepton Tagger Exclusion

Evaluated NNJQT performance with events  
not tagged by SET (Vivek's) or SMT (Gavril's)

Apply cut on **Likelihood**  $> 5\%$  for SLT (as in mixing analyses)

SLT tagged events not counted in  $\epsilon$  denominator

$D$  binning in jet type and  $|Q_{jet}|P_{nn}$

$$\epsilon_{SET} = 3.23\%$$

$$\epsilon_{SMT} = 4.41\%$$

	$\epsilon, \%$	effective $D, \%$	$\epsilon D^2$
$e + \mathbf{SVT}$	$95.12 \pm 0.15$	$8.10 \pm 0.17$	$\mathbf{0.624 \pm 0.026}$
$\mu + \mathbf{SVT}$	$95.38 \pm 0.13$	$7.98 \pm 0.15$	$\mathbf{0.607 \pm 0.023}$

Loss of 1-2% in dilution,  $\epsilon D^2$  still high

## SLT and NNJQT correlations

Measure of  $\epsilon$  and  $D$  of NNJQT on the overlap sample

No binning for NNJQT, no cuts




Likelihood  $> 5\%$  for SET and SMT

$\mu + \text{SVT}$ sample		
Subsample	$\epsilon$	$D$
SMT and NNJQT		
overlap	$4.41 \pm 0.03 \%$	$24.1 \pm 1.0 \%$
decisions agree	$3.17 \pm 0.02 \%$	$37.2 \pm 1.1 \%$
decisions disagree	$1.24 \pm 0.01 \%$	$9.74 \pm 1.8 \%$
SET and NNJQT		
overlap	$3.23 \pm 0.02 \%$	$19.2 \pm 1.1 \%$
decisions agree	$2.23 \pm 0.02 \%$	$28.0 \pm 1.3 \%$
decisions disagree	$1.00 \pm 0.01 \%$	$0.5 \pm 2.1 \%$

## Track Probability for COT only tracks


Optimize trackNet separately on  
**COT only** tracks and **SI without L00**  
 to exploit different  $d_0$  resolution

very few COT only tracks in jets (2.7%)  
 because of pre-clustering cuts on  $d_0$  and  $\Delta z_0$



## Performance of New Optimization

### Effect on jetNet



Tagging power measurement

$$0.934 \pm 0.034 \% e+SVT$$


$$0.950 \pm 0.029 \% e+SVT$$

to be compared to “standard” NNJQT

$$0.917 \pm 0.031 \% e+SVT$$

$$0.938 \pm 0.029 \% e+SVT$$

## $d_0$ and $d_0/\sigma$ contribution to trackNet



Curves for  $d_0$  and  $d_0/\sigma$  show contribution to the trackNet performance

Change of  $d_0$  and  $d_0/\sigma$  importance when COT tracks are separated is negligible

Not really necessary to optimize separately for COT only tracks



## Summary

Updated CDF note 7482, included items discussed today

Final NNJQT  $\epsilon D^2$  results are

$$\mathbf{0.917 \pm 0.031 \% \text{ } e+\textbf{SVT}}$$

$$\mathbf{0.938 \pm 0.029 \% \text{ } e+\textbf{SVT}}$$